

Practice and perception—A nationwide survey of therapy habits in sepsis*

Frank M. Brunkhorst, MD; Christoph Engel, MD; Max Ragaller, MD; Tobias Welte, MD; Rolf Rossaint, MD; Herwig Gerlach, MD; Konstantin Mayer, MD; Stefan John, MD; Frank Stuber, MD; Norbert Weiler, MD; Michael Oppert, MD; Onnen Moerer, MD; Holger Bogatsch, MD; Konrad Reinhart, MD; Markus Loeffler, MD; Christiane Hartog, MD; for the German Sepsis Competence Network (SepNet)

Objective: To simultaneously determine perceived vs. practiced adherence to recommended interventions for the treatment of severe sepsis or septic shock.

Design: One-day cross-sectional survey.

Setting: Representative sample of German intensive care units stratified by hospital size.

Patients: Adult patients with severe sepsis or septic shock.

Interventions: None.

Measurements and Main Results: Practice recommendations were selected by German Sepsis Competence Network (SepNet) investigators. External intensivists visited intensive care units randomly chosen and asked the responsible intensive care unit director how often these recommendations were used. Responses “always” and “frequently” were combined to depict perceived adherence. Thereafter patient files were audited. Three hundred sixty-six patients on 214 intensive care units fulfilled the criteria and received full support. One hundred fifty-two patients had acute lung injury or acute respiratory distress syndrome. Low-tidal volume ventilation ≤ 6 mL/kg/predicted body weight was documented in 2.6% of these patients. A total of 17.1% patients had tidal volume between 6 and 8 mL/kg predicted body weight and 80.3% > 8 mL/kg predicted body weight. Mean tidal volume

was 10.0 ± 2.4 mL/kg predicted body weight. Perceived adherence to low-tidal volume ventilation was 79.9%. Euglycemia (4.4–6.1 mmol/L) was documented in 6.2% of 355 patients. A total of 33.8% of patients had blood glucose levels ≤ 8.3 mmol/L and 66.2% were hyperglycemic (blood glucose > 8.3 mmol/L). Among 207 patients receiving insulin therapy, 1.9% were euglycemic, 20.8% had blood glucose levels ≤ 8.3 mmol/L, and 1.0% were hypoglycemic. Overall, mean maximal glucose level was 10.0 ± 3.6 mmol/L. Perceived adherence to strict glycemic control was 65.9%. Although perceived adherence to recommendations was higher in academic and larger hospitals, actual practice was not significantly influenced by hospital size or university affiliation.

Conclusions: This representative survey shows that current therapy of severe sepsis in German intensive care units complies poorly with practice recommendations. Intensive care unit directors perceive adherence to be higher than it actually is. Implementation strategies involving all intensive care unit staff are needed to overcome this gap between current evidence-based knowledge, practice, and perception. (Crit Care Med 2008; 36: 2719–2725)

KEY WORDS: sepsis; clinical practice; medical audit; questionnaires; survey

Mortality rates of severe sepsis and septic shock are still in the range of 20% to 60% (1–4). Several landmark studies have recently demonstrated

that therapeutic strategies may reduce mortality substantially. Tight glycemic control in intensive care unit (ICU) patients as a preventive approach resulted in both decreased frequency of sepsis and

reduced overall mortality (5). Lung-protective ventilation reduced absolute mortality in patients with acute respiratory distress syndrome (ARDS) by 9% (6). Early goal-directed therapy based on an algorithm including central venous O_2 saturation measurements (4), low-dose hydrocortisone in septic shock (2), and activated protein C (aPC) in appropriate patients (3) achieved further mortality reduction. On the other hand, large studies revealed that antithrombin (AT) therapy is ineffective (7) as is low-dose dopamine for presumed renal protection (8). These recommendations have been incorporated into guidelines for the treatment of severe sepsis, published by the American Society of Critical Care Medicine and other international groups (9, 10).

The gap between best practice and actual therapy habits may be associated with increased mortality and morbidity

***See also p. 2929.**

From the Anesthesiology and Intensive Care Medicine (FMB, CH, KR), Friedrich-Schiller University Jena, Germany; Institute for Medical Informatics (CE, HB, ML), Statistics and Epidemiology, University of Leipzig, Germany; Anesthesiology and Intensive Care Medicine (MR), University Hospital, Dresden, Germany; Pulmonary & Critical Care Medicine (TW), University Hannover, Germany; Anesthesiology (RR), University Hospital, Aachen, Germany; Anesthesiology and Intensive Care Medicine (HG), Vivantes-Klinikum Neukoelln, Berlin, Germany; Internal Medicine (KM), Justus Liebig University, Giessen, Germany; Department of Nephrology and Hypertension (SJ), University of Erlangen-Nuernberg, Germany; Anesthesiology and Intensive Care Medicine (FS), Rheinische Friedrich-Wilhelms-University Bonn, Germany; Anesthesiology and Intensive Care Medicine (NW), Christian-Albrechts-

University, Kiel, Germany; Nephrology and Medical Intensive Care (MO), Charité, Campus Virchow-Klinikum, Berlin, Germany; and Anesthesiology and Intensive Care Medicine (OM), University of Goettingen, Germany.

Supported in part by Federal Ministry of Education and Research, BMBF grant no. 01KI 0106, Germany and by Eli Lilly of Germany (Bad Homburg), in form of an unrestricted grant.

The authors have not disclosed any potential conflicts of interest.

For information regarding this article, E-mail: konrad.reinhart@med.uni-jena.de

Copyright © 2008 by the Society of Critical Care Medicine and Lippincott Williams & Wilkins

DOI: 10.1097/CCM.0b013e318186b6f3

(11). Only few studies have evaluated implementation of single interventions in critical care (12, 13). To date, little is known about current sepsis therapy and the uptake into clinical practice of recent major study results and recommended guidelines. Therefore, we evaluated adherence to seven recommendations in a large, nationwide, and representative sample of German ICUs.

It might be expected that larger, tertiary care institutions, when compared with smaller community hospitals, would adhere more closely to the latest study results; this has rarely been investigated. Therefore, we were careful to include a proportional selection of ICUs in hospitals from five different strata based on size.

Reports suggest that a considerable discrepancy may exist between self-assessment of guideline adherence and actual performance (14–16). Therefore, we interviewed ICU directors in a representative sample of German ICU simultaneously about their frequency of adherence to recommended best practice. This perception of adherence was compared with actual data obtained by audit of patients' medical files.

METHODS

The study was carried out by the German Sepsis Competence Network (SepNet) with 18 regional study centers, a medical coordination center (University of Jena), and a data management and biometry center (University of Leipzig). Data were collected on a cross-sectional, 1-day basis in a representative random sample of German hospitals.

Representative Hospital Sampling. A representative sample, which was stratified by size, was selected from a total of 1,380 hospitals (with a total of 488,727 beds) and 2,075 ICUs (with a total of 19,084 beds). Hospitals were divided into strata with <200, 201–400, 401–600, and >600 beds, respectively (S1–S4). Stratum 5 (S5) comprised university hospitals. A detailed description of the sampling method is given elsewhere (17).

Patients. All patients occupying an ICU bed between 6:00 AM of the study day and 6:00 AM of the following day were screened for the presence of infection, systemic inflammatory response syndrome, and organ dysfunction based on modified consensus criteria (17). Patients whose therapy was withdrawn or withheld were excluded from further analysis.

Recommended Interventions. Based on recently published recommendations by an international panel of intensivists (10) and recent major study results, the group of SepNet investigators selected seven interventions with respect to clinical relevance (Table 1). Two of

these, low-tidal volume (Vt) ventilation in patients with acute lung injury (ALI)/ARDS and glycemic control, are applicable throughout the course of the disease. Therefore, we grouped the results of the audit into time and severity-independent and time and severity-dependent recommendations.

Data Collection and Analysis. Audits were performed by external intensivists from 18 SepNet regional study centers (see Appendix) after prior central training. The auditors first gathered data on adherence to seven best practice interventions through a semistructured interview of the ICU medical director, i.e., the senior intensivist on site who is permanently in charge of local practice in the German system of closed ICUs. In the absence of the ICU director, the deputy was approached. The ICU directors could answer with “always,” “frequently,” “sometimes,” “rarely,” or “never” to each practice recommendation. Thereafter, charts of patients who fulfilled the inclusion criteria were reviewed together with the resident physician who in some cases was identical to the medical ICU director.

Patient data were anonymized in the case report forms. Electronic data entry, data management, quality control, and analysis were performed centrally. Data collected from patient documentation included maximal blood glucose levels; maximal documented Vt; administration and dosage of aPC, hydrocortisone, dopamine, and AT, all noted within 24 hrs preceding the audit.

ALI and ARDS as well as predicted body weight (PBW) were defined as described elsewhere (6). Maximum Vt (Vt max) per PBW was calculated over the preceding 24 hrs.

The study was approved by the responsible institutional ethics committees and by the federal data protection commissioner.

Statistical Analysis. We analyzed whether documented practice (i.e., maximal glucose values and Vt max in mL/kg PBW) and perceived adherence (i.e., interview responses) were significantly different across strata (i.e., hospital size). All data were categorical and are reported as absolute or relative frequencies where appropriate. Fisher's exact test was applied for all group comparisons. *p* values below 0.05 were considered significant. SPSS 11.0.1 (SPSS Inc., Chicago, IL) was used for data analysis.

RESULTS

Patients

Four hundred fifty-four ICUs from a random sample of 310 hospitals were visited, representing 22% of the total number of national ICUs. A total of 3,877 patients were screened for severe sepsis and septic shock. Four hundred fifteen patients met the criteria, representing an estimated total of 1,545 ICU patients with

Table 1. Recommended interventions (treatment goal)

Time and severity independent
Low-tidal volume ventilation ^a
(≤6 mL per kg predicted body weight)
Glycemic control ^b (blood glucose
4.4–6.1 mmol/L)
Time and severity dependent
Activated protein C ^c
Low-dose hydrocortisone
(200–300 mg/24 hr) ^d
Non-use of low-dose dopamine to protect
renal function (≤ 5 μg/kg/min) ^e
Non-use of antithrombin ^f

^aAcute lung injury as defined by Pao₂/Fio₂ >200 and <300 mm Hg or acute respiratory distress syndrome as defined by Pao₂/Fio₂ <200 mm Hg; plus bilateral infiltrates on chest x-ray and pulmonary capillary wedge pressure <18 mm Hg and/or no signs of left ventricular failure (6).

^bIntensive insulin therapy to maintain euglycemia (4.4–6.1 mmol/L) (5).

^cUsage of activated protein C (3).

^dAdministration of low-dose hydrocortisone (200–300 mg/24 hr) in patients with septic shock (2).

^eFrom Ref. (8).

^fFrom Ref. (7).

severe sepsis or septic shock on a given day in Germany. In 49 patients, therapy was withdrawn or withheld. The remaining 366 patients in 214 ICUs received full support. One hundred ninety had severe sepsis and 166 had septic shock; in a further ten patients, the presence of septic shock could not be ruled out because of incomplete documentation. Therefore, these patients were not included when patients with severe sepsis and septic shock were analyzed separately.

Intensive Care Units

One hundred eighty-seven ICUs (41.2%) were mixed surgical/medical, 85 (18.7%) were surgical, and 65 (14.3%) were medical; the median number of beds per ICU was 10 (interquartile range, 7–12). Primary specialties of ICU directors were 55.3% anesthesiology, 26.9% internal medicine, 5.7% surgery, and 11.2% others (17). ICUs were distributed among hospital strata as follows: 106 (23.3%) in S1 (≤200 beds), 151 (33.3%) in S2 (200–400 beds), 68 (15.0%) in S3 (401–600 beds), 82 (18.1%) in S4 (>600 beds), and 47 (10.4%) in university hospitals (S5).

Time and Severity-Independent Recommendations

Low-Tidal Volume Ventilation. One hundred ninety-eight patients with se-




Figure 1. Range of tidal volumes in patients with severe sepsis/septic shock and acute lung injury/acute respiratory distress syndrome. Maximal tidal volumes (mL/kg predicted body weight [PBW]) in 152 patients with severe sepsis and septic shock.




Figure 2. Range of maximal blood glucose levels in patients with severe sepsis/septic shock. Maximal blood glucose levels in 355 patients with severe sepsis and septic shock.

vere sepsis/septic shock were intubated and mechanically ventilated and fulfilled ALI/ARDS criteria. In 46 patients (23.2%), calculation of Vt in mL/kg PBW was not possible due to incomplete documentation. The remaining 152 patients were ventilated with the following Vt: four patients (2.6%) ≤ 6 mL/kg PBW, 26 patients (17.1%) between 6 and 8 mL/kg

PBW, and 122 patients (80.3%) > 8 mL/kg PBW. Mean Vt in all 152 patients was 10 ± 2.4 mL/kg PBW (Fig. 1).

Glycemic Control. Three hundred fifty-five patients with severe sepsis/septic shock had documentation of glucose values. Twenty-two of these (6.2%) were euglycemic (4.4–6.1 mmol/L), four (1.1%) had hypoglycemia (< 4.4 mmol/L). One

hundred twenty patients (33.8%) had values ≤ 8.3 mmol/L, and 235 (66.2%) had values > 8.3 mmol/L.

Two hundred seven of 355 patients were receiving insulin therapy, four of these (1.9%) were euglycemic, and two (1.0%) were hypoglycemic. Forty-three patients (20.8%) had values ≤ 8.3 mmol/L and 164 (79.2%) had values > 8.3 mmol/L. Mean maximal blood glucose in all 355 patients was 10.0 ± 3.6 mmol/L (Fig. 2).

Perceived Adherence and Comparison with Documented Practice. Neither documented practice (data not shown) nor perceived adherence to low-Vt ventilation ($p = 0.185$) or glycemic control ($p = 0.1361$) were significantly different across strata. Answers always and frequently were added to denote perceived adherence. A total of 79.9% of ICU directors perceived to adhere to low-Vt ventilation, whereas only 2.6% of patients received Vt ≤ 6 mL/kg PBW. A total of 65.9% of ICU directors perceived to adhere to glycemic control, whereas only 6.2% of patients were euglycemic.

Time and Severity-Dependent Recommendations

Activated protein C was administered to three (0.8%) of 354 patients. Low-dose hydrocortisone (200–300 mg/24 hr) was administered in 48 (30.4%) of 158 patients with septic shock and in 41 (23.3%) of 176 patients with severe sepsis. Three hundred sixteen (94.0%) of 336 patients did not receive AT, 304 (91.8%) of 331 patients did not receive low-dose dopamine. Hospital size or university affiliation had no significant effect on practiced therapy (data not shown).

Perceived Adherence. The answers of ICU directors to how often these measures were practiced in their ICUs are given in Table 2.

Based on their responses, perceived adherence was significantly different across strata with overestimation of aPC use ($p < 0.0001$) and under-estimation of low-dose dopamine use ($p < 0.0002$) and AT use ($p = 0.0421$) in larger and academic hospitals.

DISCUSSION

This study is the first to provide representative data on how glycemic control and low Vt ventilation for ALI/ARDS are used in patients with severe sepsis and septic shock in Germany.

Table 2. Perceived adherence to recommended interventions by hospital size

	All (N = 214)	S1 (n = 30)	S2 (n = 67)	S3 (n = 31)	S4 (n = 53)	S5 (n = 33)
Low-tidal volume ventilation						
Yes (%)	79.9	63.3	76.2	80.7	83.1	97.0
Not answered (%)	2.8	3.3	1.5	0.0	7.5	0.0
Glycemic control						
Yes (%)	65.9	63.3	59.7	71.0	66.1	75.8
Not answered (%)	0.5	0.0	0.0	0.0	1.9	0.0
Low-dose hydrocortisone						
Yes (%)	67.7	63.3	58.2	64.6	71.7	87.9
Not answered (%)	0.5	0.0	0.0	0.0	1.9	0.0
Activated protein C						
Yes (%)	1.4	0.0	0.0	6.5	1.9	0.0
Not answered (%)	1.4	0.0	1.5	0.0	1.9	3.0
Low-dose dopamine						
No (%)	79.0	46.7	73.2	80.6	92.5	97.0
Not answered (%)	0.5	0.0	0.0	0.0	1.9	0.0
Antithrombin						
No (%)	42.5	36.7	39.3	61.3	28.3	57.6
Not answered (%)	0.9	0.0	1.5	0.0	1.9	0.0

Auditors had a written guideline to help them conduct their interviews. They were obliged to adhere to the wording, however, were free to explain if uncertainties remained. The questions were worded as follows: "How often do you use the following measures" followed by the respective recommendation.

Response rates ranged from 97.2% to 99.5% depending on the intervention. Responses "always" and "frequently" were added to denote "yes." In case of non-use recommendations (antithrombin and low-dose dopamine), responses "never" and "rarely" were added to denote "no."

S1–S4, hospital size <200, 201–400, 401–600, and >600 beds, respectively; S5, university hospital.

Moreover, this study surveyed ICU directors' attitudes about whether recent therapy recommendations were put into practice in their ICU. The results show that while most patients did not receive recommended therapies, a majority of ICU directors responsible for these patients reported that they adhered to these recommendations.

We found that 80% of septic patients with ALI/ARDS were ventilated with Vt above 8 mL/kg PBW. These findings are in accordance with other surveys that indicate poor compliance with a strategy found to reduce mortality rates by almost 10%. Retrospective analyses of patient populations with ALI and ARDS showed mean Vt in a similar range between 10.3 ± 2.0 mL/kg PBW to 10.6 ± 2.4 mL/kg PBW (18, 19). A prospective monocenter survey using computerized prompts showed that only 39% of patients were ventilated with a Vt ≤ 7.5 mL/kg PBW 2 days after meeting ALI criteria (20). In a Finnish prospective multicentered survey in 138 ventilated patients with severe sepsis, mean Vt was 7.9 ± 2.2 mL/kg PBW. Only 15% of women were ventilated with Vt < 8 mL/kg PBW and none with less than 6 mL/kg PBW; 42.0% of men were venti-

lated with Vt < 8 mL/kg PBW and 8.2% with less than 6 mL/kg (21).

It is a matter of concern that in our study 23.2% of patients' charts lacked documentation of Vt in mL/kg. In the Finnish survey, Vt could be calculated in 138 of 247 ventilated patients (21). Other investigators have noted incomplete documentation of Vt but to a lesser degree (14). However, documentation and adequate data feedback are important for all measures of quality improvement (14, 22).

Two thirds of patients with severe sepsis or septic shock and 80% of those receiving insulin therapy were hyperglycemic. Only 6% were actually in the euglycemic range within the preceding 24 hrs. This compares with 41.5% of 470 septic patients from a recent prospective multicentered study from Finland, however, without a detailed description of the methods (21). On the other hand, hypoglycemia in our study was rare and probably due to the large proportion of patients who showed a lesser degree of glycemic control. We found that mean glucose levels were 10 mmol/L. This is higher than 8.5 mmol/L found by Krinsley (23) in a group of 800 patients on a medical-surgical ICU before implementa-

tion of an intensive glucose management protocol and substantially higher than the recommended threshold by the Surviving Sepsis Campaign (≤ 8.1 mmol/L) (24). Again, perception of practice did not match documented practice: 36% of ICU directors claimed that tight glycemic control was always used on their ICU. However, 13% decidedly stated not to use it at all (never).

Hydrocortisone use, despite high perceived acceptance (68% of ICU directors) and low treatment costs, was found in just 30% of septic shock patients, the only ones for whom it is recommended. Given the potential side effects of steroids in critically ill patients (25), it is of concern that we observed 23% of severe sepsis patients without shock who also received hydrocortisone treatment, although we cannot rule out that some of these patients had been in septic shock before and were being weaned off hydrocortisone.

There may be several reasons for non-compliance with best evidence. A considerable behavioral inertia seems to prevail despite the fact that study results and recommendations have been available in widely read journals from 1999 to 2001 (2–10, 24). This delay in knowledge translation from the time of publication of new evidence to implementation in clinical practice has been noted in other fields (11) as well as in intensive care (26). Our findings seem to rule out simple lack of awareness because interview data show that ICU directors know about and claim to practice these interventions. Various other barriers to change include doubts about the generalizability of study results (27), failure of communication especially in team work (28), differing attitudes of workers in the team (13, 29), threat to physician autonomy (30), and costs for new therapies (31). Concern for patient comfort and safety, as well as confusion regarding the diagnosis, may also affect implementation (32, 33). Each of these barriers may play a role in the interventions we studied.

Many of the recommendations of the Surviving Sepsis Campaign remain controversial. Although each of the interventions has been examined in at least one randomized controlled trial, these trials are not necessarily definitive or applicable to all patients. Therefore, residual controversy may explain some of our results. The use of steroids in septic shock remains controversial, with the recently published CORTICUS data suggesting no

benefit (34). The recommendations for glycemic control in severe sepsis are derived from a study in primarily nonseptic postsurgical patients (5) and the degree of evidence of this single-center study is considered to be low (35). A recent follow-up study in a medical ICU population, (36) as well as a recent multicentered study in patients with severe sepsis (37), did not show the same positive results as the primary study. Such concerns were also raised after the pivotal trial that resulted in the approval of aPC (38).

A striking finding is the gap between perception and clinical practice we observed. In Germany, the ICU director on the closed ICU is responsible for therapy and local practice habits. The reasons for the observed discrepancies are unclear. On one hand, this may represent a variation of the Hawthorne effect, in which the context of an investigation induces participants to improve their performance or, in this case, their perceived performance (39). On the other hand, whenever self-perception and actual practice were compared, they showed poor correlation and self-assessment ranked consistently better than performance (14–16, 29, 40). The evident mismatch between actual delivered care and physician perception of care in this setting raises concern about results obtained from observational surveys.

We observed great discrepancies between perceived and actual interventions for tight glucose control and low Vt ventilation. These require close cooperation of the entire ICU team. Poor agreement among ICU staff regarding responsibilities is a barrier to guideline implementation (13, 29). Under- or overperformance may result if attitudes and goals of care are not shared by the whole team (33, 41) whereas improvement of team communication can decrease the length of stay in the ICU (28).

To our knowledge, this is the first practice survey of ICUs spanning the full range of hospital size. Although the frequency of patients with severe sepsis increased with hospital size (17), practiced as well as perceived adherence to time and severity-independent measures were not associated with hospital size or university affiliation.

The single-point prevalence design has severe limitations regarding the conclusions from interventions which are time and severity-dependent, like hydrocortisone or AT, or require specific eligibility criteria like aPC. We also can not

rule out that patients may have received dopamine as vasopressor and were in the process of being weaned off on the day of audit. Likewise, the use of only the lowest or highest values such as the highest blood glucose value over a 24-hr period is admittedly a limited assessment of adherence to the practice of a recommended intervention. Furthermore, the audit of patient physiologic responses, which may remain out of range despite reasonable attempts to address the abnormality, is sometimes insufficient to describe the actual practice of physicians as opposed to using treatment or process measures. These findings reflect practice habits in German ICUs and may not apply to ICUs elsewhere. The strengths of the study include the high degree of representativeness covering hospitals of all sizes and a very large sample size, an interview response rate of over 98%, and external audit by trained intensivist physicians to ensure high-quality data acquisition. The patient sample represented an estimated total of 75,000 patients with severe sepsis for the year 2003 in Germany. We excluded patients for whom therapy was limited or withdrawn and thus can rule out that therapy did not comply with practice guidelines for this reason.

CONCLUSION

In this representative survey of practiced and perceived adherence to practice recommendations for the treatment of severe sepsis on German ICUs, we found that overall compliance was poor but coupled with a strong perception to the contrary in all ICUs regardless of hospital size and university affiliation. It is noteworthy that the majority of patients did not have documented low-Vt ventilation or euglycemia.

REFERENCES

1. Martin GS, Mannino DM, Eaton S, et al: The epidemiology of sepsis in the United States from 1979 through 2000. *N Engl J Med* 2003; 348:1546–1554
2. Annane D, Sebille V, Charpentier C, et al: Effect of treatment with low doses of hydrocortisone and fludrocortisone on mortality in patients with septic shock. *JAMA* 2002; 288:862–871
3. Bernard GR, Vincent JL, Laterre PF, et al: Efficacy and safety of recombinant human activated protein C for severe sepsis. *N Engl J Med* 2001; 344:699–709
4. Rivers E, Nguyen B, Havstad S, et al: Early goal-directed therapy in the treatment of se-

- vere sepsis and septic shock. *N Engl J Med* 2001; 345:1368–1377
5. van den Berghe G, Wouters P, Weekers F, et al: Intensive insulin therapy in the critically ill patients. *N Engl J Med* 2001; 345:1359–1367
6. Ventilation with lower tidal volumes as compared with traditional tidal volumes for acute lung injury and the acute respiratory distress syndrome. The acute respiratory distress syndrome network. *N Engl J Med* 2000; 342:1301–1308
7. Warren BL, Eid A, Singer P, et al: Caring for the critically ill patient. High-dose antithrombin III in severe sepsis: A randomized controlled trial. *JAMA* 2001; 286:1869–1878
8. Bellomo R, Chapman M, Finfer S, et al: Low-dose dopamine in patients with early renal dysfunction: A placebo-controlled randomised trial. Australian and New Zealand intensive care society (ANZICS) clinical trials group. *Lancet* 2000; 356:2139–2143
9. Practice parameters for hemodynamic support of sepsis in adult patients in sepsis. Task force of the American college of critical care medicine, society of critical care medicine. *Crit Care Med* 1999; 27:639–660
10. Guidelines for the management of severe sepsis and septic shock. The international sepsis forum. *Intensive Care Med* 2001; 27 (Suppl 1):S1–S134
11. Ketley D, Woods KL: Impact of clinical trials on clinical practice: Example of thrombolysis for acute myocardial infarction. *Lancet* 1993; 342:891–894
12. Kallet RH, Jasmer RM, Pittet JF, et al: Clinical implementation of the ARDS network protocol is associated with reduced hospital mortality compared with historical controls. *Crit Care Med* 2005; 33:925–929
13. Cook DJ, Meade MO, Hand LE, et al: Toward understanding evidence uptake: Semirecumbency for pneumonia prevention. *Crit Care Med* 2002; 30:1472–1477
14. Akhtar SR, Weaver J, Pierson DJ, et al: Practice variation in respiratory therapy documentation during mechanical ventilation. *Chest* 2003; 124:2275–2282
15. Pittet D, Simon A, Hugonnet S, et al: Hand hygiene among physicians: Performance, beliefs, and perceptions. *Am Intern Med* 2004; 141:1–8
16. Panu D, Lee DC: Beta-blocker use in the emergency department in patients with acute myocardial infarction undergoing primary angioplasty. *J Emerg Med* 2003; 24:379–382
17. Engel C, Brunkhorst FM, Bone HG, et al: Epidemiology of sepsis in Germany: Results from a national prospective multicenter study. *Intensive Care Med* 2007; 33:606–618
18. Thompson BT, Hayden D, Matthay MA, et al: Clinicians' approaches to mechanical ventilation in acute lung injury and ARDS. *Chest* 2001; 120:1622–1627
19. Young MP, Manning HL, Wilson DL, et al: Ventilation of patients with acute lung injury and acute respiratory distress syndrome: Has

- new evidence changed clinical practice? *Crit Care Med* 2004; 32:1260–1265
20. Kalhan R, Mikkelsen M, Dedhiya P, et al: Underuse of lung protective ventilation: Analysis of potential factors to explain physician behavior. *Crit Care Med* 2006; 34: 300–306
 21. Karlsson S, Varpula M, Ruokonen E, et al: Incidence, treatment, and outcome of severe sepsis in ICU-treated adults in Finland: The Finnsepsis study. *Intensive Care Med* 2007; 33:435–443
 22. Bradley EH, Holmboe ES, Mattern JA, et al: Data feedback efforts in quality improvement: Lessons learned from US hospitals. *Qual Saf Health Care* 2004; 13:26–31
 23. Krinsley JS: Effect of an intensive glucose management protocol on the mortality of critically ill adult patients. *Mayo Clin Proc* 2004; 79:992–1000
 24. Dellinger RP, Carlet JM, Masur H, et al: Surviving sepsis campaign guidelines for management of severe sepsis and septic shock. *Crit Care Med* 2004; 32:858–873
 25. De Jonghe B, Sharshar T, Lefaucheur JP, et al: Paresis acquired in the intensive care unit: A prospective multicenter study. *JAMA* 2002; 288:2859–2867
 26. Rubenfeld GD: Implementing effective ventilator practice at the bedside. *Curr Opin Crit Care* 2004; 10:33–39
 27. Rothwell PM: External validity of randomised controlled trials: “To whom do the results of this trial apply?” *Lancet* 2005; 365:82–93
 28. Pronovost P, Berenholtz S, Dorman T, et al: Improving communication in the ICU using daily goals. *J Crit Care* 2003; 18:71–75
 29. Slomka J, Hoffman-Hogg L, Mion LC, et al: Influence of clinicians’ values and perceptions on use of clinical practice guidelines for sedation and neuromuscular blockade in patients receiving mechanical ventilation. *Am J Crit Care* 2000; 9:412–418
 30. Reinertsen JL: Zen and the art of physician autonomy maintenance. *Ann Intern Med* 2003; 138:992–995
 31. Boldt J, Papsdorf M: [Expensive modern therapy options in intensive care medicine in Germany—Are they being used? Results of a questionnaire]. *Dtsch Med Wochenschr* 2005; 130:87–91
 32. Rubenfeld GD, Cooper C, Carter G, et al: Barriers to providing lung-protective ventilation to patients with acute lung injury. *Crit Care Med* 2004; 32:1289–1293
 33. McMullin J, Brozek J, Jaeschke R, et al: Glycemic control in the ICU: A multicenter survey. *Intensive Care Med* 2004; 30:798–803
 34. Sprung CL, Annane D, Keh D, et al: Hydrocortisone therapy for patients with septic shock. *N Engl J Med* 2008; 358:111–124
 35. Angus DC, Abraham E: Intensive insulin therapy in critical illness. *Am J Respir Crit Care Med* 2005; 172:1358–1359
 36. Van den Berghe G, Wilmer A, Hermans G, et al: Intensive insulin therapy in the medical ICU. *N Engl J Med* 2006; 354:449–461
 37. Brunkhorst FM, Engel C, Bloos F, et al: Intensive insulin therapy and pentastarch resuscitation in severe sepsis. *N Engl J Med* 2008; 358:125–139
 38. Eichacker PQ, Natanson C: Recombinant human activated protein C in sepsis: Inconsistent trial results, an unclear mechanism of action, and safety concerns resulted in labeling restrictions and the need for phase IV trials. *Crit Care Med* 2003; 31(Suppl 1): S94–S96
 39. Gale EA: The Hawthorne studies—A fable for our times? *QJM* 2004; 97:439–449
 40. O’Boyle CA, Henly SJ, Larson E: Understanding adherence to hand hygiene recommendations: The theory of planned behavior. *Am J Infect Control* 2001; 29: 352–360
 41. Thomas EJ, Sexton JB, Helmreich RL: Discrepant attitudes about teamwork among critical care nurses and physicians. *Crit Care Med* 2003; 31:956–959

APPENDIX

18 SepNet Regional Study Centers

Prof. Dr. med. R. Rossaint
 Universitätsklinikum der RWTH Aachen
 Klinik für Anästhesiologie
 Pauwelsstraße 30
 52074 Aachen

Prof. Dr. med. H. Forst / Dr. U. Jaschinski
 Klinikum Augsburg
 Klinik f. Anästhesiologie und operative Intensivmedizin
 Stanglinstr. 2
 86156 Augsburg

Prof. Dr. med. K.U. Eckardt
 Campus Virchow-Klinikum
 Med. Klinik mit Schwerpunkt Nephrologie und Intensivmedizin
 Augustenburger Platz 1
 13353 Berlin

Prof. Dr. med. H. Gerlach
 Vivantes Klinikum Neukölln
 Anästhesie und operative Intensivmedizin
 Rudower Str. 48
 12313 Berlin

PD Dr. med. F. Stüber
 Rheinische Friedrich-Wilhelms-Universität Bonn
 Klinik für Anästhesiologie und Intensivmedizin
 Sigmund-Freud-Str. 25
 53105 Bonn

PD Dr. med. M. Ragaller / Dr. K. Sengebusch
 Universität Carl Gustav Carus
 Klinik für Anästhesiologie und Intensivtherapie
 Fetscherstr. 74
 01307 Dresden

Prof. Dr. med. W. Seeger / Dr. K. Mayer
Universität Gießen
Zentrum für Innere Medizin II
Klinikstr. 29
35385 Gießen

Prof. Dr. med. H. Buchardi / Dr. O. Moerer
Georg-August-Universität Göttingen
Zentrum für Anästhesiologie, Rettungs- und Intensivmedizin
Robert-Koch-Straße 40
37075 Göttingen

Dr. med. M. Gründling
Ernst-Moritz-Arndt-Universität Greifswald
Klinik und Poliklinik für Anästhesie und Intensivmedizin
Friedrich-Löffler-Str. 23
17487 Greifswald

Prof. Dr. med. J. Radke
Martin-Luther-Universität Halle Wittenberg
Universitätsklinik für Anästhesiologie und operative Intensivmedizin
Ernst-Grube-Str. 40
06097 Halle/Saale

Prof. Dr. med. R. Brunkhorst
Klinikum Hannover-Oststadt
Klinik für Nieren-, Hochdruck- und Gefäßkrankheiten
Podbielskistr. 380
30659 Hannover

Prof. Dr. med. J. Scholz
Universitätsklinikum Kiel
Klinik für Anästhesiologie und operative Intensivmedizin
Schwanenweg 21
24105 Kiel

Dr. F.M. Brunkhorst
Klinik für Anästhesiologie und Intensivtherapie
Friedrich-Schiller-Universität Jena
Erlanger Allee 101
07740 Jena

Prof. Dr. med. D. Olthoff
Universitätsklinikum Leipzig
Klinik für Anästhesiologie und Intensivtherapie
Philip-Rosenthal-Str. 27a
04103 Leipzig

Prof. Dr. med. T. Welte
Abteilung für Pneumologie
Medizinische Hochschule Hannover
Carl-Neuberg-Str. 1
30625 Hannover

Frau Dr. med. U. Hoffmann
Klinikum Mannheim; 1. Medizinische Klinik
Theodor-Kutzer-Ufer 1-3
68167 Mannheim

PD Dr. med. H.-G. Bone
Westfälische Wilhelms-Universität Münster
Klinik und Poliklinik für Anästhesiologie und operative Intensivmedizin
Albert-Schweitzer-Str. 33
48129 Münster

PD Dr. med. S. John
Universitätsklinik Erlangen-Nürnberg
Medizinische Klinik IV
Krankenhausstr. 12
91054 Erlang
